

Anomalous ϕ Meson Suppression in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV Measured by the PHENIX Experiment at RHIC

Maxim Naglis^a for the PHENIX collaboration

^aDepartment of Particle Physics, Weizmann Institute of Science, 76100 Rehovot, Israel

Abstract

The PHENIX experiment at RHIC has measured the ϕ -meson production at mid-rapidity in $p+p$, $d+Au$, $Cu+Cu$ and $Au+Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV via the K^+K^- decay mode. The transverse momentum spectra of the ϕ -meson and the nuclear modification factor as a function of centrality are reviewed here.

1. Introduction

Quarks and gluons are believed to lose energy while traversing the strongly interacting hot and dense matter produced in relativistic heavy-ion collisions, leading to a significant softening and broadening of the jets, also known as the phenomenon of jet-quenching. Energy loss of high- p_T partons manifests itself in the suppression of the particle production at high- p_T in nucleus-nucleus collisions, compared to expectations from binary scaled $p+p$ results. The amount of suppression and its transverse momentum dependence reflect the opacity of the medium. The medium-induced effects on particle production are quantified with the nuclear modification factor:

$$R_{AA}(p_T) = \frac{d^2N_{AA}/dydp_T}{\langle N_{coll} \rangle \times d^2N_{pp}/dydp_T}, \quad (1)$$

where $d^2N_{AA}/dydp_T$ and $d^2N_{pp}/dydp_T$ are the differential yields per event in nucleus-nucleus and $p+p$ collisions, respectively, and $\langle N_{coll} \rangle$ is the number of binary nucleon-nucleon collisions averaged over the impact parameter range of the corresponding centrality.

The studies of hadron production in $p+p$, $d+Au$ and $Au+Au$ collisions in the first three runs of RHIC allowed to establish that (i) neutral pions at high- p_T are strongly suppressed in central $Au+Au$ collisions [1], (ii) no suppression is observed at high- p_T for direct photons in $Au+Au$ collisions [2] and (iii) for inclusive and identified hadrons in $d+Au$ collisions [3]. Contrary to the strong suppression of neutral pions, protons show a very different binary-collision scaling behavior from that of pions in $Au+Au$ collisions [4]. The proton production is enhanced at intermediate $p_T \approx 2-5$ GeV/ c compared to scaled $p+p$ results. The measurements by the STAR collaboration indicate that at high p_T the nuclear modification factor of protons becomes similar to that of pions [5] and also that the R_{CP} patterns for K_S^0 and $K(892)^*$ mesons are different from that of the Λ baryon (note that $m_{K(892)^*} \approx m_\Lambda$) [6, 7]. Further studies of identified hadron production reveal that despite a factor of about 4 difference in mass, η mesons follow the suppression pattern of neutral pions over the entire p_T range of the measurements [8]. The latter suggests that the differences in the behavior of baryons and mesons are not related to the difference in

their mass but rather to the number of constituent quarks. The ϕ -meson, with a mass comparable to that of the proton and $\Lambda(1115)$ baryon, but carrying two quarks, differentiates between hadron mass and number of constituent quark effects. Moreover, being an almost pure $s\bar{s}$ state, it provides insight on the effects of flavour composition on the hadron suppression pattern.

2. Measurements of the $\phi \rightarrow K^+ K^-$ with the PHENIX Detector

The $\phi \rightarrow K^+ K^-$ results presented in this contribution were obtained from the data samples accumulated by the PHENIX experiment [9] during $p+p$, $d+\text{Au}$, $\text{Cu}+\text{Cu}$ and $\text{Au}+\text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV in the 2003-2005 physics runs. Charged particle tracking and measurements of their momentum are accomplished by the high-resolution multi-wire proportional Drift Chambers (DC) and the first layer of the Pad Chambers (PC1). The typical value of the momentum resolution is $\sigma(p_T)/p_T \approx 1.0\% p_T \oplus 1.1\%$. Charged particle identification (PID) is based on the particle mass calculated from the momentum and the time-of-flight information derived from the Time of Flight (TOF) detector or the Lead Scintillator (PbSc) part of the Electro Magnetic Calorimeter and the Beam Beam Counters (BBC's). The TOF subsystem with a time resolution $\sigma \approx 120$ ps and the PbSc with $\sigma \approx 500$ ps allow to achieve reliable pion-kaon separation in the p_T ranges $0.3 - 2.5$ GeV/ c and $0.3 - 1.0$ GeV/ c , respectively. The Zero Degree Calorimeters (ZDC's) and BBC's are dedicated subsystems that determine the collision vertex and event centrality and also provide the minimum bias (MB) interaction trigger. The MB trigger used for $p+p$, $d+\text{Au}$ and $\text{Cu}+\text{Cu}$ collisions requires a coincidence between the BBC's with at least one hit in each BBC arm. For $\text{Au}+\text{Au}$ collisions the MB trigger requires a coincidence between the BBC's and ZDC's with at least two hits in each BBC arm, and at least one neutron detected in each ZDC arm. All events used in the analysis are required to have the collision vertex position along the beam axis within 30 cm of the geometrical center of PHENIX. There are three different techniques that were used for the $\phi \rightarrow K^+ K^-$ mass reconstruction. The first ("no PID") does not require identification of charged tracks in the final state and assumes that all tracks are kaons. The second ("one kaon PID") requires identification of only one kaon in the TOF. In the third technique ("two kaons PID") both kaons are identified in the TOF or the PbSc; four different subsystem combinations have been considered: TOF-TOF, TOF-PbSc_{EAST}, PbSc_{EAST}-PbSc_{EAST}, PbSc_{WEST}-PbSc_{WEST}.

The $d+\text{Au}$ data were analyzed using "no PID", $p+p$ and $\text{Cu}+\text{Cu}$ using "no PID" and "one kaon PID", and $\text{Au}+\text{Au}$ using "no PID" and "two kaons PID". Combining the results obtained with complementary techniques the p_T range accessible for the ϕ measurements extends from ~ 1 GeV/ c to 7 GeV/ c (to 5.1 GeV/ c in $d+\text{Au}$). The three analysis techniques have very different sources of systematic uncertainties and provide a valuable consistency check. In every technique all kaon candidates from each event passing the track selection requirements are combined into unlike-sign pairs. The invariant mass and the transverse momentum for each pair is then calculated based on the 2-body decay kinematics. The resulting mass distributions contain both the ϕ -meson signal and an inherent combinatorial background. The combinatorial background is estimated either by simultaneously fitting the mass distribution to the sum of a Breit-Wigner function to account for the signal, convoluted with Gaussian to account for the mass resolution, and a polynomial function to account for the background, or by an event-mixing technique described elsewhere [10]. The raw yields are extracted by integrating the invariant mass distributions in the vicinity of the PDG value of the ϕ -meson mass (1.019 GeV/ c^2) after subtracting the combinatorial background. Corrections to the raw yields for the limited detector acceptance and

resolution, reconstruction and trigger efficiency, multiplicity effects and various analysis cuts are determined from the full single-particle Monte Carlo simulation and analysis of the data.

3. Results




Figure 1: Invariant p_T spectra of the ϕ -meson measured in (left) $p+p$, $d+Au$, $Au+Au$ and (right) $Cu+Cu$ collisions at $\sqrt{s_{NN}} = 200$ GeV by the PHENIX experiment.

The ϕ -meson invariant p_T spectra measured in $p+p$, $d+Au$, $Cu+Cu$ and $Au+Au$ collisions are summarized in Fig. 1. Left panel of the figure shows good agreement between the various techniques. Using the combined $p+p$ results as a reference, the nuclear modification factor has been derived for the ϕ -meson in $d+Au$, $Au+Au$ and $Cu+Cu$ collisions. In minimum bias and most central $d+Au$ collisions, the modification factor R_{dA} shows no suppression of ϕ -mesons [11]. Fig. 2 shows the R_{AA} for the ϕ in the most central $Au+Au$ collisions over a p_T range of 2.45-7 GeV/c. There is an ongoing work to extend this p_T range towards low- p_T . The R_{AA} for π^0 , $(K^+ + K^-)/2$, η , ω , $(p + \bar{p})/2$, and direct γ are also shown in Fig.2 for comparison. High- p_T direct photons show no suppression up to $p_T \approx 14$ GeV/c [12]. Protons are enhanced at $p_T > 1.5$ GeV/c [13]. Neutral pions and η mesons follow the same suppression pattern [8]. ϕ -mesons appear to be less suppressed compared to π^0 and η mesons in the p_T range of $2.45 < p_T < 4.5$ GeV/c. At higher p_T (> 5 GeV/c) the amount of suppression of ϕ and ω mesons could be similar to that of π^0 and η , unfortunately both statistical and systematic errors are too large for a more conclusive statement. The similarity between the suppression patterns of different mesons at high- p_T supports the concept of high- p_T particle production via the fragmentation of partons outside the hot and dense medium. It is not clear whether or not the R_{AA} for the kaon (also containing a strange quark) follows the trend of the ϕ , since the present measurements have no overlap in p_T .

Comparative studies of $Au+Au$ and $Cu+Cu$ data sets allow to test how sensitive is the particle production at high- p_T to the collision geometry. The ϕ 's R_{AA} measurements performed in $Au+Au$ and $Cu+Cu$ collisions for similar number of participating nucleons $\langle N_{part} \rangle$, i.e. similar energy density, yield similar results. Fig. 3 shows no significant difference in the suppression patterns for 40%-50% central $Au+Au$ ($\langle N_{part} \rangle = 74.4$) and 10%-20% central $Cu+Cu$ ($\langle N_{part} \rangle = 73.6$) collisions in the overlapping p_T range. This is illustrated more generally in Fig. 4 where the integrated nuclear modification factor is shown as a function of $\langle N_{part} \rangle$. Despite the differences in the amount of suppression between the ϕ and π^0 , good agreement in the suppression levels of the same meson is observed in $Au+Au$ and $Cu+Cu$ collisions for similar $\langle N_{part} \rangle$ [14].




Figure 2: Nuclear modification factor R_{AA} in central Au+Au collisions as a function of p_T for π^0 , $(K^+ + K^-)/2$, η , ω , $(p + \bar{p})/2$, ϕ and direct γ .




Figure 3: Comparison of the ϕ 's R_{AA} measured in Au+Au and Cu+Cu collisions for similar number of participating nucleons $\langle N_{part} \rangle$.

4. Summary

The PHENIX experiment has measured the ϕ -meson production in $p+p$, $d+Au$, Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV via the K^+K^- decay channel. The suppression pattern for the ϕ -meson measured in central Au+Au collisions is drastically different at intermediate- p_T from that for π^0 and η mesons. The ϕ 's R_{AA} for Au+Au and Cu+Cu collisions shows a similar dependence on p_T and $\langle N_{part} \rangle$.

Acknowledgments

The author acknowledges support by the Israel Science Foundation, the MINERVA Foundation and the Nella and Leon Benoziyo Center of High Energy Physics Research.

References

- [1] S. Adler et al., *Phys. Rev. Lett.* **91** (2003) 072301.
- [2] S. Adler et al., *Phys. Rev. Lett.* **94** (2005) 232301.
- [3] S. Adler et al., *Phys. Rev. Lett.* **91** (2003) 072303.
- [4] S. Adler et al., *Phys. Rev. Lett.* **91** (2003) 172301.
- [5] B. Abelev et al., *Phys. Rev. Lett.* **91** (2006) 152301.
- [6] J. Adams et al., *Phys. Rev. Lett.* **92** (2004) 052302.
- [7] J. Adams et al., *Phys. Rev.* **C71** (2005) 064902.
- [8] B. Sahlmueller, *J. Phys.* **G34** (2007) S969-974.
- [9] K. Adcox et al., *Nucl. Instrum. Methods* **A499** (2003) 469.
- [10] S. Adler et al., *Phys. Rev.* **C72** (2005) 014903.
- [11] V. Ryabov, *J. Phys.* **G35** (2008) 044030.
- [12] K. Reygers, *J. Phys.* **G35** (2008) 104045.
- [13] S. Adler, *Phys. Rev.* **C74** (2006) 024904.
- [14] H. Buesching, *Eur. Phys. J.* **C49** (2006) 41.




Figure 4: ϕ -meson's integrated R_{AA} for Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV.